## Statistics Homework 2: Probability

## Javier Tasso

- 1. Solve the following exercises of chapter 2 of Devore, Berk, Carlton's *Modern Mathematical Statistics with Applications* (third edition): 1, 3, 4, 9, 11, 13, 14, 16, 19, 23, 31, 45, 48, 49, 50, 51, 57, 69, 70, 71, 74, 75, 78, 79, 81, 117, 127
- 2. Prove the following statement: if  $A \subset B$  then P(A) < P(B).
- 3. It is well known that  $P(A \cup B) = P(A) + P(B) P(A \cap B)$ . Use this result to show:

$$P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(A \cap C) - P(B \cap C) + P(A \cap B \cap C)$$

- 4. Explain intuitively why  $\binom{n}{k}$  equals  $\binom{n}{n-k}$ .
- 5. Suppose P(B) > 0 show that P(A|B) + P(A'|B) = 1.
- 6. Prove the following statements:
  - (a) If A and B are independent, then A and B' are independent too.
  - (b) If A and B are independent, then A' and B' are independent too.

## Answers

- 1. You can find answers to the odd numbered problems at the end of the textbook.
  - 4. a.  $A = \{RRR, LLL, SSS\}$ 
    - b.  $B = \{RLS, RSL, LSR, LRS, SLR, SRL\}$
    - c.  $C = \{RRL, RRS, RLR, RSR, SRR, LRR\}$
    - d.  $D = C \cup \{LLS, LLR, LSL, LRL, RLL, SLL\} \cup \{SSL, SSR, SRS, SLS, RSS, LSS\}$
    - e.  $D' = A \cup B$ ,  $C \cup D = D$ ,  $C \cap D = C$ .
  - 14. a. 0.65
    - b. 0.35
    - c.  $A \cap B'$ . The probability is 0.25
  - 16. a. 0.31
    - b. 0.69
    - c. 0.37
  - 48. a. 2520 by doing  $\binom{10}{2} \cdot \binom{8}{3} \cdot \binom{5}{5}$ .
    - b. Simplify in the previous expression:  $\frac{10! \cdot 8! \cdot 5!}{2!8! \cdot 3!5! \cdot 5!0!} = \frac{10!}{2!3!5!}$ .
    - c. n! in the numerator counts every possible permutation of n objects, treating all arrangements as distinct. The denominator  $k_1! \ldots k_r!$  corrects for over-counting due to the fact that the internal arrangement within each group does not matter.
  - 50. You should expect P(A|B) to be higher. The universe of professional basketball players is much smaller than the universe of tall people.
  - 70. Let S be the event that the student is satisfied with the textbook. By the law of total probability P(S) = 0.51. Now update to get the posterior probabilities P(Mean|S) = 0.3922, P(Median|S) = 0.2941, P(Mode|S) = 0.3137. It is still more likely he used professor Mean's textbook.
  - 74. P(A) = 0.5 and P(A|B) = 0.625. These are different, so A and B are not independent. You can verify that  $P(A) \cdot P(B)$  does not equal  $P(A \cap B)$ .
  - 78.  $P(OO) = 0.44 \cdot 0.44 = 0.1936$  and P(Match) = P(AA) + P(BB) + P(ABAB) + P(OO) = 0.3816
- 2. A is contained in B. You can use a Venn diagram to verify that  $B = A \cup (B \cap A')$  where A and  $(B \cap A')$  are disjoint. Then:

$$B = A \cup (B \cap A')$$

$$P(B) = P[A \cup (B \cap A')]$$

$$P(B) = P(A) + P(B \cap A')$$

Now notice that  $P(B \cap A') > 0$ . Then P(B) > P(A).

3. Define  $D = B \cup C$ 

$$P(A \cup D) = P(A) + P(D) - P(A \cap D)$$

$$= P(A) + P(B \cup C) - P[A \cap (B \cup C)]$$

$$= P(A) + P(B) + P(C) - P(B \cap C) - P[A \cap (B \cup C)]$$

$$= P(A) + P(B) + P(C) - P(B \cap C) - P[(A \cap B) \cup (A \cap C)]$$

$$= P(A) + P(B) + P(C) - P(B \cap C) - [P(A \cap B) + P(A \cap C) - P(A \cap B \cap C)]$$

Rearranging you get the desired formula.

- 4. Suppose you have n = 10 people and you want to count the number of ways you can create groups of k = 8. By selecting groups of 8 you are also selecting groups of 2. This is way the two combinations match. If you select a group of 8 people, the remaining 2 are automatically chosen in the unselected group.
- 5. Recall that  $P(B) = P(A \cap B) + P(A' \cap B)$ . Divide both sides by P(B), use the definition of the conditional probability.
- 6. (a) Start with  $P(A) = P(A \cap B) + P(A \cap B')$ , use independence, and rearrange.

$$P(A) = P(A \cap B) + P(A \cap B')$$

$$P(A) = P(A)P(B) + P(A \cap B')$$

$$P(A) - P(A)P(B) = P(A \cap B')$$

$$P(A)[1 - P(B)] = P(A \cap B')$$

$$P(A)P(B') = P(A \cap B')$$

The last equality shows that A and B' are independent.

(b) Focus on B' and A. From the previous property we know they are independent (because A and B were).

$$P(B') = P(A \cap B') + P(A' \cap B')$$

$$P(B') = P(A)P(B') + P(A' \cap B')$$

$$P(B') - P(A)P(B') = P(A' \cap B')$$

$$P(B')[1 - P(A)] = P(A' \cap B')$$

$$P(B')P(A') = P(A' \cap B')$$